

FILE 'REGISTRY' ENTERED AT 15:18:33 ON 03 SEP 2008

L1 STRUCTURE UPLOADED

L2 0 S L1


L3 6 S L1 SSS FULL

FILE 'HCAPLUS' ENTERED AT 15:19:39 ON 03 SEP 2008

L4 6 S L3

=>

Uploading C:\Program Files\STNEXP\Queries\10546132markush.str



chain nodes :

7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27
28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48
49 50 51 52
53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73
74 76 78 85
87

ring nodes :

1 2 3 4 5 6

chain bonds :

1-10 1-69 2-11 2-68 3-7 3-72 5-9 5-71 6-8 6-70 7-76 8-85 9-87 10-47
11-78 12-13 12-45 13-14 14-15 14-46 15-16 16-17 17-18 18-19 19-20 20-21
21-22 22-23
23-24 24-25 26-27 26-44 27-28 28-29 29-30 30-31 31-32 32-33 33-34 34-35
35-36 36-37 36-73
37-38 37-74 38-39 39-40 40-41 41-42 42-43 47-48 48-49 49-50 49-57 49-58
50-51 51-52
52-53 53-54 54-55 55-56 59-60 59-66 60-61 60-65 62-63 63-64 64-67

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6
 exact/norm bonds :
 1-2 1-6 1-10 2-3 2-11 3-4 4-5 5-6 5-9 6-8 7-76 8-85 9-87 11-78 12-45
 14-46 26-44 49-57 63-64
 exact bonds :
 1-69 2-68 3-7 3-72 5-71 6-70 10-47 12-13 13-14 14-15 15-16 16-17 17-18
 18-19 19-20 20-21 21-22 22-23 23-24 24-25 26-27 27-28 28-29 29-30 30-31
 31-32 32-33
 33-34 34-35 35-36 36-37 36-73 37-38 37-74 38-39 39-40 40-41 41-42 42-43
 47-48 48-49
 49-50 49-58 50-51 51-52 52-53 53-54 54-55 55-56 59-60 59-66 60-61 60-65
 62-63 64-67

G1:H, Si, CH3

G2:H, P

G3:H, MeO

G4:[*1], [*2]

G5:[*3], [*4]

Match level :
 1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS 8:CLASS 9:CLASS 10:CLASS
 11:CLASS 12:CLASS 13:CLASS 14:CLASS 15:CLASS 16:CLASS 17:CLASS 18:CLASS
 19:CLASS 20:CLASS
 21:CLASS 22:CLASS 23:CLASS 24:CLASS 25:CLASS 26:CLASS 27:CLASS 28:CLASS
 29:CLASS 30:CLASS
 31:CLASS 32:CLASS 33:CLASS 34:CLASS 35:CLASS 36:CLASS 37:CLASS 38:CLASS
 39:CLASS 40:CLASS
 41:CLASS 42:CLASS 43:CLASS 44:CLASS 45:CLASS 46:CLASS 47:CLASS 48:CLASS
 49:CLASS 50:CLASS
 51:CLASS 52:CLASS 53:CLASS 54:CLASS 55:CLASS 56:CLASS 57:CLASS 58:CLASS
 59:CLASS 60:CLASS
 61:CLASS 62:CLASS 63:CLASS 64:CLASS 65:CLASS 66:CLASS 67:CLASS 68:CLASS
 69:CLASS
 70:CLASS 71:CLASS 72:CLASS 73:CLASS 74:CLASS 76:CLASS 78:CLASS 85:CLASS
 87:CLASS

L1 STRUCTURE UPLOADED

=> d l1

L1 HAS NO ANSWERS

L1 STR

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

Structure attributes must be viewed using STN Express query preparation.

=> s l1

SAMPLE SEARCH INITIATED 15:19:11 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 85 TO ITERATE

100.0% PROCESSED

85 ITERATIONS

0 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 1147 TO 2253
PROJECTED ANSWERS: 0 TO 0

L2 0 SEA SSS SAM L1

=> s l1 sss full

FULL SEARCH INITIATED 15:19:18 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 2031 TO ITERATE

100.0% PROCESSED 2031 ITERATIONS
SEARCH TIME: 00.00.01


6 ANSWERS

L3 6 SEA SSS FUL L1

=> d l3 scan

L3 6 ANSWERS REGISTRY COPYRIGHT 2008 ACS on STN
IN α -D-Glucopyranoside, (1Z)-1-propen-1-yl 3-O-decyl-2-deoxy-2-[(1,3-dioxotetradecyl)amino]-
MF C33 H61 N O7

Absolute stereochemistry.
Double bond geometry as shown.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):2

L3 6 ANSWERS REGISTRY COPYRIGHT 2008 ACS on STN
IN α -D-Glucopyranoside, (1Z)-1-propen-1-yl 3-O-decyl-2-deoxy-6-O-[(1,1-dimethylethyl)dimethylsilyl]-2-[(1,3-dioxotetradecyl)amino]-
MF C39 H75 N O7 Si


Absolute stereochemistry.
Double bond geometry as shown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L3 6 ANSWERS REGISTRY COPYRIGHT 2008 ACS on STN
 IN β -D-Glucopyranoside, (1E)-1-propen-1-yl 3-O-decyl-2-deoxy-6-O-[(1,1-dimethylethyl)dimethylsilyl]-2-[(1,3-dioxotetradecyl)amino]-
 MF C39 H75 N O7 Si

Absolute stereochemistry.
 Double bond geometry as shown.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L3 6 ANSWERS REGISTRY COPYRIGHT 2008 ACS on STN
 IN α -D-Glucopyranoside, (1Z)-1-propen-1-yl 2-deoxy-3-O-[(3R)-3-methoxydecyl]-6-O-methyl-2-[(11Z)-1-oxo-11-octadecen-1-yl]amino]-
 MF C39 H73 N O7

Absolute stereochemistry.
 Double bond geometry as shown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):0

=> file hcaplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

178.82

179.03

FILE 'HCAPLUS' ENTERED AT 15:19:39 ON 03 SEP 2008

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 3 Sep 2008 VOL 149 ISS 10

FILE LAST UPDATED: 2 Sep 2008 (20080902/ED)

HCAPLUS now includes complete International Patent Classification (IPC) reclassification data for the second quarter of 2008.

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s l3

L4 6 L3

=> d l4 1-6 ti abs bib hitstr

L4 ANSWER 1 OF 6 HCAPLUS COPYRIGHT 2008 ACS on STN

TI Sodium salt of glucosamine disaccharide compound, method for producing the same, and use of the same

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB There are disclosed a sodium salt represented by the average formula (I; m1, n1, m2 and n2 independently represent 0 or a pos. number not more than 2, while satisfying $m1 + n1 = 2$, $m2 + n2 = 2$, $0 < m1 + m2 < 4$ and $0 < n1 + n2 < 4$.) and a method for producing such a sodium salt. There is also a decomposition suppressing method which enables to have a sodium salt represented by the average formula I coexistent with a sodium salt represented by the general formula II below. This method enables to improve long-term stability of a sodium salt represented by the general formula II which is effective for the prevention and/or treatment of septicemia caused by gram pos. bacteria, in particular endotoxin shock. Thus, a DEAE column main fraction containing 6.0 g disaccharide free acid (III) (preparation given) and

4.80

weight% Na and 942.8 L MeOH were stirred in a 4 L flask at 25°, treated with 0.2 N NaOH/MeOH solution (15.2 mL), stirred overnight, filtered, and treated dropwise with 270 mL acetone at 25°. The precipitate was removed by filtration and dried in vacuo to give III.3.67 Na. When III.3.67 Na was stored in a screw-cap bottle at 25° for 30 days, impurities A, B, and C were formed at a rate of 0.072, 0.267, and 0.072 %/mo, resp., vs. 0.729, 3.117, and 0.033 %/mo, resp., for III.4.06Na.

AN 2008:636616 HCAPLUS <<LOGINID::20080903>>

DN 149:10241

TI Sodium salt of glucosamine disaccharide compound, method for producing the same, and use of the same

IN Sakurai, Shin; Furukawa, Ken; Matsuo, Kimihiro; Tagami, Kenichi

PA Eisai R & D Management Co., Ltd., Japan

SO PCT Int. Appl., 46pp.

CODEN: PIXXD2

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	---	-----	-----	-----
PI	WO 2008062842	A1	20080529	WO 2007-JP72579	20071121
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW				
	RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				

PRAI JP 2006-315020 A 20061122

US 2006-860483P P 20061122

IT 748165-18-6P 748165-20-0P

RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)


(intermediate; preparation of sodium salt of glucosamine disaccharide compound

with storage stability, method for producing it, and its use for prevention and/or treatment of endotoxin shock)

RN 748165-18-6 HCAPLUS

CN α -D-Glucopyranoside, (1Z)-1-propen-1-yl 2-deoxy-3-O-[(3R)-3-methoxydecyl]-6-O-methyl-2-[[(11Z)-1-oxo-11-octadecen-1-yl]amino]-, 4-(di-2-propen-1-yl phosphate) (CA INDEX NAME)


Absolute stereochemistry.
Double bond geometry as shown.



RN 748165-20-0 HCAPLUS

CN α -D-Glucopyranose, 2-deoxy-3-O-[(3R)-3-methoxydecyl]-6-O-methyl-2-[[(11Z)-1-oxo-11-octadecen-1-yl]amino]-, 4-(di-2-propen-1-yl phosphate) 1-(2,2,2-trichloroethanimidate) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.



IT 748165-17-5

RL: RCT (Reactant); RACT (Reactant or reagent)


(preparation of sodium salt of glucosamine disaccharide compound with storage

stability, method for producing it, and its use for prevention and/or treatment of endotoxin shock)

RN 748165-17-5 HCAPLUS


CN α -D-Glucopyranoside, (1Z)-1-propen-1-yl 2-deoxy-3-O-[(3R)-3-methoxydecyl]-6-O-methyl-2-[[(11Z)-1-oxo-11-octadecen-1-yl]amino]- (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.



RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 6 HCAPLUS COPYRIGHT 2008 ACS on STN
TI Process for production of lipid A analogue
GI



I

AB There is disclosed a process for producing 3-O-decyl-2-deoxy-6-O-[2-deoxy-3-O-[(3R)-3-methoxydecyl]-6-O-methyl-2-[[11Z]-1-oxo-11-octadecenyl]amino]-4-O-phosphono-β-D-glucopyranosyl]-2-[(1,3-dioxotetradecyl)amino]-α-D-glucopyranose 1-(dihydrogen phosphate) (known as eritoran) tetrasodium salt (I; R = PO₃Na₂) which is useful as an active ingredient of a pharmaceutical or an intermediate for the synthesis thereof. A process for producing the compound I (R = PO₃Na₂) comprises the key steps of reacting a compound represented by the formula I [R = P(O)(OCH₂CH:CH₂)₂] with a palladium catalyst in the presence of a nucleophilic agent (deallylation) and treating the product with a sodium source (sodium salt formation). This process is environment-friendly and excellent in safety, operability, and reproducibility. Thus, a solution of 101.6 g I [R = P(O)(OCH₂CH:CH₂)₂] in 203 mL THF was added to a mixture of Meldrum's acid 70.49, palladium acetate 2.93, and PPh₃ 51.3 g and the resulting mixture was stirred at 32° for 2 h and at 30° for 4 h, treated with 250 mL MeOH, and concentrated under reduced pressure to give a residue (466.7 g). The residue was dissolved in 4,570 mL MeOH at 40°, treated with

5.55 g trimercaptotriazine, stirred overnight at room temperature, and filtered to remove the precipitated trimercaptotriazine-palladium complex, followed by washing the precipitate with MeOH to give a combined filtrate (4,330 g). The filtrate (3,908.2 mL) was concentrated under reduced pressure to give a residue (440.9 g) which was treated with 450 mL acetone, concentrated under reduced pressure, treated again with 450 mL acetone, and concentrated under reduced pressure. The residue was refrigerated overnight, treated with 1,800 mL acetone, warmed to 40°, stirred for 1.5 h, air-cooled, stirred at ≥30° for 1.5 h, and filtered to give, after washing with acetone and drying at 35-40° under reduced pressure, 74.2% eritoran (free acid form) which was treated with 0.1 N aqueous NaOH solution to give eritoran tetrasodium salt.

AN 2007:257680 HCAPLUS <<LOGINID::20080903>>

DN 146:317153

TI Process for production of lipid A analogue

IN Tagami, Katsuya; Sato, Keizo; Matsuo, Kimihiro; Abe, Taichi; Haga, Toyokazu

PA Eisai R & D Management Co., Ltd., Japan

SO PCT Int. Appl., 69pp.

CODEN: PIXXD2


DT Patent

LA Japanese

FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2007026675	A1	20070308	WO 2006-JP316941	20060829
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW				
	RW:				
	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	AU 2006285926	A1	20070308	AU 2006-285926	20060829
	CA 2620027	A1	20070308	CA 2006-2620027	20060829
	EP 1939209	A1	20080702	EP 2006-796921	20060829
	R:				
	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, RS				
	KR 2008039374	A	20080507	KR 2008-700572	20080109
	CN 101238140	A	20080806	CN 2006-80027144	20080124
PRAI	US 2005-712431P	P	20050831		
	JP 2005-253044	A	20050901		
	WO 2006-JP316941	W	20060829		
OS	CASREACT 146:317153				
IT	748165-18-6P 748165-20-0P				
	RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)				
	(process for production of lipid A analog (eritoran) via palladium-catalyzed deallylation of eritoran diallyl ester and formation of sodium salt)				
RN	748165-18-6 HCAPLUS				
CN	α-D-Glucopyranoside, (1Z)-1-propen-1-yl 2-deoxy-3-O-[(3R)-3-methoxydecyl]-6-O-methyl-2-[[[(11Z)-1-oxo-11-octadecen-1-yl]amino]-, 4-(di-2-propen-1-yl phosphate) (CA INDEX NAME)				

Absolute stereochemistry.
Double bond geometry as shown.





RN 748165-20-0 HCAPLUS
 CN α -D-Glucopyranose, 2-deoxy-3-O-[(3R)-3-methoxydecyl]-6-O-methyl-2-
 [[(11Z)-1-oxo-11-octadecen-1-yl]amino]-, 4-(di-2-propen-1-yl phosphate)
 1-(2,2,2-trichloroethanimidate) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.



IT 748165-17-5
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (process for production of lipid A analog (eritoran) via
 palladium-catalyzed deallylation of eritoran diallyl ester and
 formation of sodium salt)
 RN 748165-17-5 HCAPLUS
 CN α -D-Glucopyranoside, (1Z)-1-propen-1-yl 2-deoxy-3-O-[(3R)-3-
 methoxydecyl]-6-O-methyl-2-[[(11Z)-1-oxo-11-octadecen-1-yl]amino]- (CA
 INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.




FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2007039450	A	20070215	JP 2006-187298	20060707
PRAI	JP 2005-199518	A	20050708		

OS MARPAT 146:252059
 IT 859508-28-4P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (preparation of glucose lipid analogs for treatment of inflammation and
 autoimmune diseases)
 RN 859508-28-4 HCAPLUS
 CN β -D-Glucopyranoside, (1E)-1-propen-1-yl 3-O-decyl-2-deoxy-6-O-[(1,1-
 dimethylethyl)dimethylsilyl]-2-[(1,3-dioxotetradecyl)amino]- (CA INDEX
 NAME)


Absolute stereochemistry.
 Double bond geometry as shown.



L4 ANSWER 4 OF 6 HCAPLUS COPYRIGHT 2008 ACS on STN
 TI Syntheses of glucose-containing E5564 analogs and their LPS-antagonistic
 activities
 AB Lipid A analogs containing a glucose moiety on their non-reducing end were
 synthesized, and their LPS-antagonistic activities were measured. The
 inhibitory activities (IC₅₀) on LPS-induced TNF α production of title
 aminodeoxy disaccharides toward human whole blood cells were 0.46-1.11 nM.
 Inhibitory doses (ID₅₀) of these compds. on TNF α production induced by
 co-injection of galactosamine and LPS in C3H/HeN mice were measured. The
 ID₅₀ values of these compds. were 0.20-1.08 and <0.2 mg/kg. Moreover,
 C3H/HeN mice preinjected with compds. were protected from lethality
 induced by co-injection of galactosamine and LPS. Out of eight mice
 preinjected with 1 mg/kg of title compds. five-eight mice were protected.
 AN 2005:1299295 HCAPLUS <<LOGINID::20080903>>
 DN 144:171174
 TI Syntheses of glucose-containing E5564 analogs and their LPS-antagonistic
 activities
 AU Shiozaki, Masao; Doi, Hiromi; Tanaka, Daisuke; Shimozato, Takaichi;
 Kurakata, Shin-ichi
 CS Chemistry Department, Chemtech Labo, Inc., Hiromachi 1-2-58, Shinagawa-ku,
 Tokyo, 140-8710, Japan
 SO Tetrahedron (2005), Volume Date 2006, 62(1), 205-225
 CODEN: TETRAB; ISSN: 0040-4020
 PB Elsevier B.V.
 DT Journal
 LA English
 OS CASREACT 144:171174
 IT 859508-28-4P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (syntheses of glucose-containing E5564 analogs and their LPS-antagonistic
 activities)


RN 859508-28-4 HCAPLUS
 CN β -D-Glucopyranoside, (1E)-1-propen-1-yl 3-O-decyl-2-deoxy-6-O-[(1,1-dimethylethyl)dimethylsilyl]-2-[(1,3-dioxotetradecyl)amino]- (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry as shown.



RE.CNT 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 5 OF 6 HCAPLUS COPYRIGHT 2008 ACS on STN
 TI preparation of levulose glucoselipid A derivatives as TNF α
 production inhibitors
 GI



AB Title compds. I [Q = O, etc.; W = O, NH; R1 = (un)substituted alkanoyl, etc. with the proviso that if W = NH; R1 (with the proviso that if W = O), R2, R3, R4 = H, (un)substituted alkyl, etc.; R5 = H, halo, etc.] were prepared For example, phosphorylation of 4-O-(allyloxycarbonyl)-3-O-decyl-2-deoxy-6-O-[4-O-diallylphosphono-3-O-[(R)-3-methoxydecyl]-6-O-methyl-2-O-[(Z)-11-octadecenoyl]- β -D-glucopyranosyl]-2-(3-oxotetradecanoylamino)- α -D-glucopyranoside, e.g., prepared from 1,2:5,6-di-O-isopropylidene- α -D-glucofuranose in 15 steps, with diallyl diisopropylphosphoramidate followed by deallylation using Pd(PPh₃)₄ afforded phosphono 3-O-decyl-2-deoxy-6-O-[3-O-[(R)-3-methoxydecyl]-6-O-methyl-2-O-[(Z)-11-octadecenoyl]-4-O-phosphono- β -D-glucopyranosyl]-2-(3-oxotetradecanoylamino)- α -D-glucopyranoside (II). In TNF α production inhibition assays, the IC₅₀ value of compound II was 0.49 nM. Compds. I are claimed useful for the treatment of inflammation, septicemia, etc.

AN 2005:638895 HCAPLUS <<LOGINID::20080903>>

DN 143:153644

TI preparation of levulose glucoselipid A derivatives as TNF α production inhibitors

IN Shiozaki, Masao; Shimozato, Takaichi

PA Sankyo Company, Limited, Japan

SO PCT Int. Appl., 156 pp.

CODEN: PIXXD2

DT Patent


LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005066193	A1	20050721	WO 2005-JP434	20050107
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2552218	A1	20050721	CA 2005-2552218	20050107
	JP 2005220130	A	20050818	JP 2005-2028	20050107
	EP 1702926	A1	20060920	EP 2005-703673	20050107
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK, IS				
	CN 1930180	A	20070314	CN 2005-80007429	20050107
	BR 2005006671	A	20070515	BR 2005-6671	20050107
	IN 2006KN01892	A	20070511	IN 2006-KN1892	20060706
	MX 2006PA07822	A	20060926	MX 2006-PA7822	20060707
PRAI	JP 2004-2902	A	20040108		
	WO 2005-JP434	W	20050107		
OS	MARPAT 143:153644				
IT	859508-28-4P				
	RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)				
	(preparation of levulose glucoselipid A derivs. as TNF α production inhibitors for treatment of inflammation, septicemia, etc.)				
RN	859508-28-4 HCAPLUS				
CN	β -D-Glucopyranoside, (1E)-1-propen-1-yl 3-O-decyl-2-deoxy-6-O-[(1,1-dimethylethyl)dimethylsilyl]-2-[(1,3-dioxotetradecyl)amino]- (CA INDEX NAME)				

Absolute stereochemistry.

Double bond geometry as shown.



RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 6 OF 6 HCAPLUS COPYRIGHT 2008 ACS on STN
TI Reagents and methods for preparing lipopolysaccharides antagonist B1287
and stereoisomers thereof for treatment of various forms of septic shock
GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The present invention provides methods for preparing lipopolysaccharides (LPS) antagonist lipo-disaccharide B1287 and stereoisomers thereof, which compds. are useful as in the prophylactic and affirmative treatment of endotoxemia including sepsis, septicemia and various forms of septic shock (no biol. data). Also provided are synthetic intermediates useful for implementing the inventive methods. Thus, lipo-disaccharide B1287 I was prepared for treatment of various forms of septic shock.

AN 2004:718552 HCAPLUS <<LOGINID::20080903>>

DN 141:225771

TI Reagents and methods for preparing lipopolysaccharides antagonist B1287 and stereoisomers thereof for treatment of various forms of septic shock

IN Fan, Rulin

PA Eisai Co, Ltd., Japan

SO PCT Int. Appl., 175 pp.

CODEN: PIXXD2

DT Patent


LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004074303	A2	20040902	WO 2004-US4921	20040218
	WO 2004074303	A3	20041229		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI			
	RW:	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	JP 2006518394	T	20060810	JP 2006-503710	20040218
	US 20060160999	A1	20060720	US 2005-546132	20051212
PRAI	US 2003-448839P	P	20030220		
	WO 2004-US4921	W	20040218		
OS	CASREACT 141:225771; MARPAT 141:225771				
IT	748165-17-5P 748165-18-6P 748165-20-0P 748165-24-4P 748165-25-5P				
	RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)				
	(reagents and methods for preparing lipopolysaccharides antagonist b and stereoisomers thereof for treatment of various forms of septic shock)				
RN	748165-17-5 HCAPLUS				
CN	α -D-Glucopyranoside, (1Z)-1-propen-1-yl 2-deoxy-3-O-[(3R)-3-methoxydecyl]-6-O-methyl-2-[[[(11Z)-1-oxo-11-octadecen-1-yl]amino]- (CA INDEX NAME)				

Absolute stereochemistry.

Double bond geometry as shown.




RN 748165-18-6 HCAPLUS

CN α -D-Glucopyranoside, (1Z)-1-propen-1-yl 2-deoxy-3-O-[(3R)-3-methoxydecyl]-6-O-methyl-2-[[[(11Z)-1-oxo-11-octadecen-1-yl]amino]-, 4-(di-2-propen-1-yl phosphate) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry as shown.




RN 748165-20-0 HCAPLUS

CN α -D-Glucopyranose, 2-deoxy-3-O-[(3R)-3-methoxydecyl]-6-O-methyl-2-[[[(11Z)-1-oxo-11-octadecen-1-yl]amino]-, 4-(di-2-propen-1-yl phosphate) 1-(2,2,2-trichloroethanimidate) (CA INDEX NAME)

Absolute stereochemistry.


Double bond geometry as shown.



RN 748165-24-4 HCAPLUS

CN α -D-Glucopyranoside, (1Z)-1-propen-1-yl 3-O-decyl-2-deoxy-2-[(1,3-dioxotetradecyl)amino]- (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.



RN 748165-25-5 HCAPLUS

CN α -D-Glucopyranoside, (1Z)-1-propen-1-yl 3-O-decyl-2-deoxy-6-O-[(1,1-dimethylethyl)dimethylsilyl]-2-[(1,3-dioxotetradecyl)amino]- (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.

